

400

**Forecast skill of targeted
observations: a singular vector
based diagnostic**

C. Cardinali and R. Buizza

Research Department

February 2003

Accepted by the Journal of Atmospheric Sciences

For additional copies contact:For additional copies please contact

The Library
ECMWF
Shinfield Park
Reading, Berks RG2 9AX

library@ecmwf.int

Series: ECMWF Technical Memoranda

A full list of ECMWF Publications can be found on our web site under: <http://www.ecmwf.int/publications.html>

© Copyright 2003

European Centre for Medium Range Weather Forecasts
Shinfield Park, Reading, Berkshire RG2 9AX, England

Literary and scientific copyrights belong to ECMWF and are reserved in all countries. This publication is not to be reprinted or translated in whole or in part without the written permission of the Director. Appropriate non-commercial use will normally be granted under the condition that reference is made to ECMWF.

The information within this publication is given in good faith and considered to be true, but ECMWF accepts no liability for error, omission and for loss or damage arising from its use.



Summary

Targeted dropsonde data have been assimilated using the operational ECMWF 4D-Var system for 10 cases of the NORth Pacific EXperiment (NORPEX) campaign, and their impact on analyses and corresponding forecasts has been investigated. The 10 fastest growing “analysis” singular vectors have been used to define a sub-space of the phase space where initial conditions are expected to be modified by the assimilation of targeted observing. A linear combination of this vector basis is the pseudo-inverse, that is the smallest perturbation with the largest impact on the forecast error. The dropsonde-induced analysis difference has been decomposed into three initial perturbations, two belonging to the sub-space spanned by the leading 10 SVs and one to its complement. Differences and similarities of the three analysis components have been examined, and their impact on the forecast error compared with the impact of the pseudo-inverse.

Results show that, on average, the dropsonde-induced analysis difference component in the sub-space spanned by the leading 10 SVs and the dropsonde-induced analysis difference component along the pseudo-inverse directions are very small (6% and 15%, respectively, in terms of total energy norm). In the only case where dropsonde data were exactly released in the area identified by the SVs, the different components of the dropsonde-induced analysis difference and the pseudo-inverse had consistent impacts on the forecast error. It is concluded that the poor agreement between the dropsonde location and the SV maxima is the main reason for the relatively small impact of the NORPEX targeting observations on the forecast error.

1. Introduction

Rapidly developing cyclones that form towards the end of the Atlantic and Pacific storm-tracks are sometimes difficult to forecast. The sparsity of observational data over the oceans can result in analysis errors which may grow rapidly in the ensuing forecast. Following the first ideas discussed at a workshop in 1995 (*Snyder 1996*), several field experiments have been carried out to observe atmospheric circulations in traditionally data sparse regions and to assess whether the assimilation of extra observations in a target area can improve forecast quality in a downstream verification area. Field experiments include the Fronts and Atlantic Storm Track Experiment (FASTEX, *Thorpe and Shapiro 1995*), the Pacific Experiment (NORPEX, *Langland et al 1999*), the California Landfalling Jets Experiment (CALJET, *Emanuel et al 1995, Ralph et al 1998*) and the Winter Storm Reconnaissance Experiment (WSR99, *Szunyogh et al 2000*, and WSR00, *Szunyogh et al 2001*). Results based on the 18 cases from the Winter Storm Reconnaissance programs (*Szunogh et al 1999, Toth et al 2002*), for example, indicated forecast improvement in 60-70% of the cases, during which the surface pressure root-mean-square errors inside a preselected verification areas have been measured to decrease by 10%. Similarly, results based on 4 FASTEX cases (*Montani et al 1999*) reported a 15% average decrease of the root-mean-square forecast error for the 500 and 1000 hPa geopotential height fields.

One of the key problems is that it is not obvious where best to deploy the dropsonde data. Several approaches to identifying the sensitive regions have been proposed and used in targeting campaigns: the Sensitivity Vectors (*Rabier et al. 1996, Langland et al. 1996 and 1999, Gelaro et al. 1998*), the Ensemble Transform Technique (ETT, *Bishop and Toth 1999*) and the Singular Vector (SV) technique (*Buizza and Montani 1999; Gelaro et al. 1999*). The reader is referred to published literature (e.g. *Palmer et al., 1998*) for a discussion of similarities and differences among these techniques. Targeting techniques also include the Quasi-Inverse linear method (*Pu et al., 1997 and 1999*) and the Ensemble Transform Kalman Filter (ETKF, *Bishop et al., 2001*, the ETKF had been used operationally as targeting guidance during the 2000, 2001 and 2002 WSR missions).

This study explores the forecast impact results from the assimilation of targeting dropsondes during the Pacific EXperiment (NORPEX), one of the first experiments designed to investigate the possible benefits of real time targeting. NORPEX took place in mid-January and February 1998 with dropsondes deployed by NOAA and U.S. Air Force aircraft to improve 1-3 day forecasts over the west coast of United States. Two targeting techniques were used during NORPEX: the first one based on the ETKF implemented at NCEP and the second one based on SVs computed at NRL. In the ETKF an index of analysis sensitivity computed from an ensemble of forecasts determines the target locations, while in the SV technique, target locations are defined by an index based on the weighted average of the initial-time SVs computed to maximize total energy inside the verification area.

The comparison between the two types of forecast, one starting from analysis generated with targeted dropsondes and one without, indicates that targeting was successful only in 7 out of 10 NORPEX cases (see Section 3). The fact that targeting does not always reduce the forecast error may be due to many possible reasons: a wrong definition of the target area, an inconsistency between the assimilation procedure and the definition of the target area (one of the weaknesses of the “energy norm” SV targeting technique is that it does not take into account the characteristics of the data assimilation

system used to assimilate targeted observations), a non-optimal assimilation of the targeted observation, and model errors.

This paper reports results from data-assimilation and forecast experiments designed to investigate possible reasons of the small positive impact on the forecast error obtained for the selected NORPEX cases. SVs are used only as a diagnostic tool to investigate the impact on the forecast error of targeted dropsonde data and, thus, the strengths and weaknesses of a targeting technique based on SVs are not discussed, nor the efficiency of the ETKF and SV targeting techniques is compared. After this introduction, Section 2 describes the NORPEX campaign and the SV-based diagnostic technique. Results from analysis and forecast experiments are discussed in Section 3 and 4. Conclusions are drawn in section 5.

2. Targeting in data-sparse mid-latitude regions

2.1 The NORPEX campaign

In Winter 1997-1998, heavy precipitation occurred over parts of California, probably associated with maximum intensity of El Niño towards the end of January. During this period, the atmospheric circulation was dominated by a strong jet-level wind with storms releasing large amounts of rain over the California coast. One of the primary goals of the NORPEX campaign was to improve the short range forecast in a specified forecast verification area (FVA) of the - Western American Coast. During the 27 days of the NORPEX experiment, 3 NOAA and 2 US Air Force aircraft released almost 700 dropsondes over the eastern Pacific, with a horizontal separation of 100 to 250 km. The dropsondes provided vertical profiles of temperature, wind, humidity and pressure from the aircraft level (300-400 hPa) to the surface. These observations were mainly released at 00 UTC and were distributed in real time via the Global Telecommunication Service network to meteorological centres.

Some inconsistencies were found in the humidity values discouraging their use in the analyses (*Jaubert et al.* 1999), and thus only wind and temperature measurements have been assimilated. In the assimilation, the same observation error as far radiosondes is assigned to the dropsondes. Targeted observations were released in areas identified by NRL and NCEP using two different techniques: NRL targets were defined using the first 4 SVs computed with the NOGAPS model with a fixed verification area (FVA, 30-60N, 100-130W) and with a 2-day optimization time interval (*Langland et al.* 1999), while NCEP targets were defined applying the ETKF technique to NCEP and ECMWF global ensemble forecasts, with a flow-dependent verification area and with variable lead times (1 to 2 days; *Toth et al.* 1999, *Szunyogh et al.* 2000). At ECMWF, data from 10 NORPEX campaigns with initial states at 00UTC of 7, 9, 11, 15, 18, 20, 22, 25, 26 and 27 of February 1998 were received. These ten cases, chronologically numbered from 1 to 10, are investigated in this work.

2.2 Singular vector based diagnostics

SVs identify perturbations with the fastest growth during a finite time interval, called the optimization time interval. SVs can grow from the analysis time $t=0$ to the optimization time interval (“analysis SVs”), or from an initial to a final forecast (“forecast SVs”). Either type of SVs forms an orthogonal basis at the initial and final times with respect to the chosen metric. The Appendix briefly summarizes the SV mathematical definition.

At ECMWF, “analysis” SVs have been used routinely to define the initial perturbations of the Ensemble Prediction System (EPS, *Buizza and Palmer* 1995, *Molteni et al.* 1996). The rationale of this choice was that the analysis error component along the leading SVs dominates the forecast error. At the time of writing (September 2002), the SVs used to define the perturbed analysis of the ECMWF EPS are computed with a T42L40 resolution (spectral triangular truncation T42 with 40 vertical levels) and with a 48 hour optimization time interval.

The use of the SVs in targeting applications is the natural extension of published results (*Buizza et al.*, 1997, *Gelaro et al.*, 1998) showing that the correction of the initial conditions with a perturbation defined by the leading SVs can significantly improve the forecast skill inside a chosen FVA. More specifically, a linear combination of the leading SVs can be used to define the pseudo-inverse initial perturbation that can correct the most of the forecast error inside the FVA. The pseudo-inverse is computed from the forecast error projection onto the evolved SVs inside the FVA (Appendix A). The forecast error reduction induced by the pseudo-inverse initial perturbation can be used as an upper bound forecast error reduction that can be achieved by adding a small perturbation to the initial conditions (*Buizza and Montani* 1999). For this reason, the pseudo-inverse initial perturbation is used as a reference initial perturbation in this study.

It should be pointed out that during real-time targeting experiments “forecast” SVs growing from 24 or 36 hour forecast are used instead of “analysis” SVs to allow a sufficient lead-time for flight preparation. By contrast, “analysis” SVs are

used in this study as a diagnostic tool for an a-posteriori assessment of the impact of targeted observations in 10 NORPEX cases. As shown by *Gelaro et al. (1999)*, “analysis” SVs are more appropriate than “forecast” SVs for diagnosing forecast behavior and investigating possible reasons of success and failure of targeting experiments. The reader is also referred to *Buizza and Montani (1999)* for a discussion on the similarities among “analysis” and “forecast” SVs computed with different lead times.

The “analysis” SVs (hereafter simply called SVs) have been computed with a T63L31 model version with simplified dry (i.e. without any moist process included in the tangent and adjoint model version) physics (*Buizza 1994*), with a 48-hour optimization time period and with the final-time total energy optimized inside a fixed FVA defined by the coordinates: 30-60N and 100-130W (that is the area used by NRL during the real-time experiment). Note that the same resolution T63L31 is used in the 4D-Var data-assimilation experiments to compute the analysis increments (see Section 3).

Computer resources limit to few tens the number of SVs that can be routinely computed in a reasonable amount of time. Figure 1 shows the percentage of forecast error that projects onto a different number of leading SVs. Results show that this percentage is fast increasing up to 44% (on average) for the first 10 vectors, whilst it increases only by another 3% (on average) by adding the next 10 leading SVs. This suggests that using 10 SVs is a good compromise between a realistic representation of the fast-growing error component and the high SV computational cost. It is worth mentioning that during the NORPEX campaign only 4 SVs were used to identify sensitive regions (*Langland et al. 1999*).




Figure 1: Percentage of the forecast error that projects onto the leading evolved SVs as a function of the number of singular vectors, averaged over the 10 NORPEX cases.

SV growth is measured using a total energy norm, which is the most commonly used metric in predictability studies (*Palmer et al., 1998*). As a consequence, the SVs (and the pseudo-inverse) are computed without any knowledge of the characteristics of the data assimilation system used to generate the analysis (e.g. observation and background covariance matrices). Thus, there is no guarantee that the pseudo-inverse perturbation is similar to the modification induced by the assimilation of the extra targeted observations. In the adaptive observation problem, information about the analysis errors distribution can have a significant impact on targeting location and optimal sampling of the observations. *Ehrendorfer and Tribbia (1997)* suggested a way to link SV structures and data-assimilation system by using as initial norm an estimate of the analysis errors covariance matrix (Hessian). *Barkmeijer et al (1999)* computed Hessian SVs and compared their characteristics with the routinely computed total energy SVs. Their results did not show any significant change in the percentage of forecast error explained by the leading SVs, however, the structure can be different (*Leutbecher et al. 2002*).

In this study, the leading 10 SVs define the sub-space where initial conditions are expected to be modified by the assimilation of dropsonde observations. The net effect of assimilating dropsonde data is represented by the difference between the analyses computed with and without them. Hereafter, this difference is named the dropsonde-induced analysis difference. In order to understand the role of targeted observations on the forecast error, the relationships between the dropsonde-induced analysis difference, the SV sub-space and the pseudo-inverse have been investigated.

3. Methodology: sv-based diagnostics and forecast error

Two data-assimilation experiments have been performed: a control experiment, experiment *C*, with all the observations operationally used at ECMWF in which no dropsonde data were assimilated, and experiment *D* with NORPEX dropsonde observations also included. Both experiments used the ECMWF 4D-Var data-assimilation system (*Rabier et al. 2000, Mahfouf and Rabier 2000, Klinker et al. 2000*) in a configuration with a T319L31 (T319 spectral triangular truncation with 31 vertical levels) high-resolution model integrations with full physical parametrization and T63L31 low-resolution minimizations with simplified physics (*Mahfouf 1999*). During the assimilation, the high-resolution 6-hour forecast is compared with all available observations over a six hour period whilst the analysis increments are computed at a T63L31 resolution. The *C* and the *D* analyses have been generated via a continuous data-assimilation, and 2 day forecast have been performed for the 10 NORPEX cases.

For any NORPEX case, let f^c and f^d be the 48 hour forecasts started from the initial analyses a_0^c and a_0^d , respectively, and let a^c and a^d be the *C* and *D* analyses verifying the 48 hour forecasts ($t=0$ is the targeting time). The *C* and *D* forecast errors are given by:

$$\begin{aligned} e^c &= f^c - a^c \\ e^d &= f^d - a^c \end{aligned} \quad (1)$$

with both forecast errors computed with respect to the *C* analysis since

$$\|da\| = \|a^c - a^d\| \ll \|a^c\| \quad (2)$$

and the forecast error is little sensitive to using either of the two verifying analyses. $\|\cdot\|$ denotes the total energy norm (see Appendix A).

Define δa as the T63 truncation of the analysis difference $da = a_0^c - a_0^d$ inside the area T centred on the region where the dropsondes were released (Pacific, 20-60N, 140-240E), expressed in terms of upper-air vorticity, divergence and temperature, and surface pressure components. The T63 spectral truncation and the exclusion of the specific humidity components make δa comparable to the SVs and the pseudo-inverse. The geographical restriction to the area T guarantees that, for each case, the dropsonde-induced analysis perturbation δa is mostly determined by the dropsondes released on that precise day. Results discussed in the following sections will indicate that approximating da with δa has a negligible impact on forecast error evolution inside the FVA in 9 out of the 10 cases.

Three different initial perturbations have been defined by decomposing the analysis perturbation δa in two different ways to allow the forecast error impact investigation of the dropsonde-induced analysis difference and its relationship with the leading SVs and the pseudo-inverse initial perturbation. The first initial perturbation has been defined by decomposing δa as

$$\delta a = \delta a_{\parallel} + \delta a_{\perp} \quad (3)$$

where δa_{\parallel} and δa_{\perp} are the components parallel and orthogonal to the phase-space direction defined by the pseudo-inverse initial perturbation δp (see Appendix A). δa_{\parallel} defines the first initial perturbation.

The other two initial perturbations have been defined by decomposing δa as

$$\delta a = \delta a_{SV} + \delta a_{\overline{SV}} \quad (4)$$

where δa_{SV} is the projection of δa onto the sub-space defined by the leading 10 SVs and $\delta a_{\overline{SV}}$ is its projection onto the complement sub-space. Note that, since the SVs and the pseudo-inverse perturbation are defined only in terms of the model's upper-level vorticity, divergence, temperature and surface pressure, the same applies to δa , δa_{\parallel} , δa_{SV} and $\delta a_{\overline{SV}}$.

The pseudo-inverse δp , the three initial perturbations δa_{\parallel} , δa_{SV} , $\delta a_{\overline{SV}}$ and δa have been subtracted from the *C* analysis to define five perturbed initial conditions:

$$\begin{aligned}
 a_0^1 &= a_0^c - \delta p \\
 a_0^2 &= a_0^c - \delta a_{\parallel} \\
 a_0^3 &= a_0^c - \delta a_{SV} \\
 a_0^{\bar{3}} &= a_0^c - \delta a_{\overline{SV}} \\
 a_0^4 &= a_0^c - \delta a
 \end{aligned} \tag{5}$$

from each of which forecast experiments (number similarly) have been run to assess the impact of each perturbation on the forecast error in the FVA:

- forecasts f^1 and f^c are compared to assess the impact of the pseudo-inverse perturbation;
- forecasts f^2 and f^c are compared to assess the impact of the δa component along the pseudo-inverse;
- forecasts f^3 , $f^{\bar{3}}$ and f^c are compared to assess the impact of the δa component that belongs to the sub-space defined by the leading 10 SVs and its complement;
- forecasts f^4 and f^c are compared to assess the impact of the dropsonde-induced perturbation δa . Forecast results are discussed in terms of the relative forecast error

$$RE(f^j) = \frac{\|e^j\| - \|e^c\|}{\|e^c\|} \tag{6}$$

where each forecast error $\|e^j\|$ is measured by the square root of the total energy norm inside the FVA (vertically integrated). The relative forecast error $RE()$ is the change (in percentage) of the forecast error with respect to the control forecast: a negative $RE(f^j)$ indicates that f^j has a smaller error than the control forecast.

3.1 Impact on the forecast error of the initial perturbation δa and role of specific humidity

First, the impact of approximating da by δa is investigated. Figure 2 shows the relative forecast error of the forecast f^d started from the D analysis ($a_0^d = a_0^c - da$) and of the forecast f^4 started from the control analysis plus the truncated and localized dropsonde-induced analysis perturbation ($a_0^4 = a_0^c - \delta a$). Note that the difference between $RE(f^4)$ and $RE(f^d)$ is very small (smaller than 0.02 for 5 cases and between 0.02 and 0.04 for 4 cases) for all but one case, case number 3, for which the difference is 0.15. Figure 2 also indicates that the time evolution of the T63 upper-air vorticity, divergence and temperature and surface pressure components of the dropsonde-induced analysis difference are dominant with respect to the small-scales ($>T63$) and to the specific humidity component. This is not surprising since in this study the analysis




Figure 2: Normalized forecast error $RE()$ with error measured in terms of a vertically integrated total-energy norm, and normalized by the control forecast error and averaged over the FVA (Eq. (6)) for the 10 NORPEX cases. Black column, $RE(f^d)$, white column, $RE(f^4)$ and grey column, $RE(f^q)$ show the impact of different components of the drop-induced analysis increments: superscript (d) states for full drop-induced analysis increments, (4) for the drop-induced analysis increments inside T at T63 resolution and (q) as (4) but with the humidity field of D analysis (details in par. 3).

increments are computed at T63 resolution, while the higher T319 resolution is used only when the model trajectory and the observations are compared at the observation point.

Another perturbed analysis, a_0^q , has been defined to investigate whether neglecting the specific humidity in δa is the main reason for the difference between $RE(f^4)$ and $RE(f^d)$ in case 3. The analysis a_0^q is defined by replacing the humidity field of a_0^4 with the a_0^d humidity field (i.e. a_0^q includes δa and the humidity analysis perturbation induced by the assimilation of the dropsonde data). Figure 2 shows that $RE(f^q)$ is very similar to $RE(f^d)$, with differences smaller than 2% for all cases including case 3, suggesting that the difference between $RE(f^4)$ and $RE(f^d)$ for this case is indeed due to the lack of humidity component in δa . The fact that the assimilation of temperature and wind profiles from targeted observations can induce changes in the specific humidity field is not surprising. In fact, although dropsondes specific humidity is not directly assimilated at ECMWF, mass and wind observations can generate humidity increments due to the dynamical link between temperature and humidity induced by the virtual temperature and by the action of the simplified linearized physics used in the minimization.

3.2 Impact on the forecast error of the initial perturbations δa_{SV} and $\delta a_{\overline{SV}}$

Initial perturbations δa_{SV} and $\delta a_{\overline{SV}}$ have been defined in Section 3 to investigate the relationship between the dropsonde-induced analysis perturbation and the SVs. The following ψ index

$$\psi = \frac{\|\delta a_{SV}\|}{\|\delta a\|} \quad (7)$$

can be used to measure the relative amplitude of the analysis component projecting onto the leading 10 SVs.

Table 1 shows that on average $\psi \approx 6\%$ with peak value of $\psi = 9\%$ for cases 1 and 6. This indicates, that the projection of the dropsonde-induced difference onto the sub-space defined by the leading ten SVs is small (less than one tenth of the total analysis difference).




Figure 3: Normalized forecast error $RE()$ with error measured in terms of a vertically integrated total-energy norm, and normalized by the control forecast error and averaged over the FVA (Eq. (6)). Black column, $RE(f^4)$, white column, $RE(f^{\bar{3}})$ and grey column $RE(f^3)$ show the impact of different components of the drop-induced analysis increments: superscript (4) states for the drop-induced analysis increments inside T at T63 resolution, (3) for the drop-induced analysis increments projecting onto the SVs sub-space and ($\bar{3}$) for its complement (details in par. 3).

Table 1: Amplitude of δa_{SV} relative to the dropsonde-induced perturbation

Case	<u>1</u>	2	3	4	5	<u>6</u>	7	8	9	10
$\psi(\%)$	<u>9</u>	5.5	4.6	6.5	6.4	9	6.3	2.7	4.8	4

Figure 3 shows the relative forecast error $RE(f^4)$ (started from $a_0^4 = a_0^c - \delta a$), $RE(f^3)$ (started from $a_0^3 = a_0^c - \delta a_{SV}$) and $RE(f^3)$ (started from $a_0^3 = a_0^c - \delta a_{SV}$). In five cases (2, 4, 5, 8 and 10) $RE(f^4) - RE(f^3)$ which indicates the δa_{SV} component of the dropsonde-induced analysis perturbation determines the impact of δa on the forecast error. For three other cases (3, 6, and 9) $RE(f^4) \sim RE(f^3)$, i.e. the impact of δa on the forecast error is determined by δa_{SV} , while for the last two cases (1 and 7) both components have a comparable contribution.

Overall, these results indicate that, the component of the dropsonde-induced perturbation along the leading 10 SVs, δa_{SV} dominates the forecast evolution only in 3 cases, while the complement perturbation $\delta a_{\perp SV}$ dominates in 5 cases. Moreover, the relative forecast impact modulo ($|RE(f^3)|$) is not much related to the amplitude ψ ; one would have expected larger $|RE(f^3)|$ for larger ψ , but only case 6, with maximum amplitude, shows the largest forecast impact over the 10 cases. Impact on the forecast error of the pseudo-inverse δp and δa_{\parallel}

Figure 4 shows the relative forecast error $RE(f^4)$ (started from $a_0^4 = a_0^c - \delta a$), $RE(f^1)$ (started from $a_0^1 = a_0^c - \delta p$) and $RE(f^2)$ (started from $a_0^2 = a_0^c - \delta a_{\parallel}$) for the 10 campaigns. $RE(f^1) < 0$ indicates that the pseudo-inverse δp always induces a forecast error reduction, and being $RE(f^1) < RE(f^4)$ indicates that the pseudo-inverse δp always corrects the forecast error more than δa . The fact that $RE(f^1) < 0$ is qualitatively in agreement with the result expected if the pseudo-inverse time evolution was linear, but it should be pointed out that there is a disagreement between the average forecast error reduction $\langle RE(f^1) \rangle = 10\%$ (Fig. 4) and the forecast error projection onto the leading 10 SVs that is on average 44% (Fig. 1). This discrepancy indicates that non-linear processes have an important impact on the time evolution of the pseudo-inverse. Other possible reasons for this disagreement can rely on the simplified physical processes described in the tangent and adjoint model version used to compute the SVs (e.g. moist processes and radiation are not included) (Buizza and Montani 1999, Gilmour et al 2001).




Figure 4: Normalized forecast error $RE()$ with error measured in terms of a vertically integrated total-energy norm, and normalized by the control forecast error and averaged over the FVA (Eq. (6)). Black column, $RE(f^4)$, white column, $RE(f^1)$ and grey column $RE(f^2)$ show the impact of different components of the drop-induced analysis increments: superscript (4) states for the drop-induced analysis increments inside T at T63 resolution, (1) for pseudo-inverse perturbation and (2) for the drop-induced analysis increments projecting onto pseudo-inverse (details in par. 3).

To quantify the relationship between the pseudo-inverse δp and the analysis difference δa two other indices have been defined. The first index ρ is the ratio between δa_{\parallel} and the norm of δp

$$\rho = \frac{\langle \delta a, \frac{\delta p}{\|\delta p\|} \rangle}{\|\delta p\|} \quad (8)$$

where $\langle \cdot, \cdot \rangle$ denotes the inner product defined in terms of the total energy norm. Positive (negative) ρ values indicate that δa_{\parallel} points along the same (opposite) direction as the pseudo-inverse. If $\rho=1$, then δa_{\parallel} has the same amplitude as the pseudo-inverse perturbation. The second index is the angle α between the two vectors δp and δa

$$\alpha = \arccos \frac{\langle \delta a, \delta p \rangle}{\|\delta p\| \|\delta a\|} \quad (9)$$

Table 2 shows that ρ is smaller than 0.1 and α is close

Table 2: Amplitude and angle between the vectors δa and δp

Case	1	2	3	4	5	<u>6</u>	<u>7</u>	8	9	10
ρ	0.05	-0.07	-0.28	0.14	-0.05	<u>0.60</u>	<u>0.58</u>	0.08	-0.01	0.04
α°	86°	100°	113°	61°	99°	<u>38°</u>	<u>44°</u>	65°	92°	81°

to 90° for all but four cases (3, 4, 6 and 7): cases 6 and 7 which have $|\rho|>0.5$, and cases 3 and 4 which have $0.5>|\rho|>0.1$. Table 2 indicates that the dropsonde-induced analysis difference δa has a small component along the pseudo-inverse and almost perpendicular direction. In other words, the dropsonde-induced difference and the pseudo-inverse perturbation are similar in 2 cases (number 6 and 7) and different or very different in the other 8 cases.

Figure 4 shows the impact on the forecast error of $\delta a_{||}$ and δp . Consider first the four cases with $|\rho|>0.1$ and smaller α (cases 3, 4, 6 and 7). Results show that for cases 6 and 7, characterized by the largest positive ρ ($\rho=0.60$ and 0.58 , respectively), $RE(f^2)\sim RE(f^1)$. For case number 4 ($\rho=0.14$), $RE(f^2)\sim 0.2*RE(f^1)$, while for case number 3 ($\rho=-0.28$), $RE(f^2)$ is about three-times smaller and has the opposite sign of $RE(f^1)$. For the other 6 cases characterized by $|\rho|<0.1$ there is no correspondence between ρ and the forecast error impact of $\delta a_{||}$ and δp .

Despite the fact that clear cut conclusions cannot be drawn from this set of results, the indication is that δa and δp have a similar impact on the forecast error when a large enough fraction of the dropsonde-induced analysis difference δa projects onto the pseudo-inverse δp , say when $\rho>0.58$ (2 out of 10 cases). Results also show that there is still a certain degree of agreement when $0.14<|\rho|<0.58$ (2 out of 10 cases), but that no relationship can be found when $|\rho|<0.1$.

3.3 Dropsonde location

The results discussed in the previous sections have indicated that the dropsonde-induced analysis difference has a small component on the sub-space spanned by the leading SVs, and that, on average, there is a very little agreement between the dropsonde induced analysis difference and the pseudo-inverse. One possible reason of this disagreements could be that the dropsondes were released in areas that did not coincide with the area of maximum concentration of the (“analysis”) SVs used in this study both to define δa_{SV} and δp . The “analysis” SV which sample a very similar area as the “forecast” SV (compare Fig 10 in this paper with Fig 5 of *Majumdar et al.*, 2002), can be used to map the general location of maximum sensitivity of the real-time leading SVs.




Figure 5: Modulo of the relative forecast error $|RE(f^3)|$ versus DLE (Dropsonde Location Efficiency, see text for definition). Labels indicate the NORPEX campaign number and the regression $Y=0.63X+0.008$ is the solid line.

The agreement between the locations of maximum SV concentration and the dropsonde has been quantified by the *DLE* (Dropsonde Location Efficiency) index defined by the sum of the SV energy (weighted mean of total energy) at the observations locations divided by the sum of the SV energy over the area T. Large *DLE* values indicate that grid points

with high average SV concentration are sampled ($DLE=1$ if the dropsondes sampled the whole area identified by the leading SVs, and $DLE=0$ if the dropsondes have been released outside the area sampled by the SVs). Figure 5 shows a scatter plot of the moduli of $RE(f^3)$ as a function of DLE . The moduli of $RE(f^3)$ are strongly correlated with DLE . In fact, although the small sample size, the high correlation found (0.81) is significantly different from zero (p-value less than 0.01). The regression line has a significant positive slope, 0.63, while the intercept is not significant (p-value=0.14). In conclusion, the data show a clear relationship. On average, dropsondes sample 2.3% of the area of maximum concentration identified by the “analysis” SVs, and in the most successful campaign case number 6 (Fig 5), DLE has a maximum value of 8%. From the scatter plot it can be seen that the impact of δa_{SV} is large for cases with large DLE .




Figure 6: Case 5, initial state 00GMT of the 18th of February 1998, 500 geopotential height fields. a) δa perturbation, contours every 2 m. b) Difference between the 2-day forecast absolute-error of f^4 and f^c started from a_0^4 and C analyses and valid on the 20 February 1998 at 00 UTC. Contour every 6 m. c) δa_{sv} perturbation, contours every 0.4 m. d) Difference between the 2-day forecast absolute-error of f^3 and f^c , contour every 6 m. e) Difference between the 2-day forecast absolute-error of f^3 and f^c , contour every 6 m. Shaded contours are negative.

4. Case studies

A detailed discussion of two cases is reported hereafter to give the reader a visual and more complete picture of the relationship between the dropsonde-induced perturbation δa , the pseudo-inverse δp and the three defined initial perturbations δa_{SV} , $\delta a_{\overline{SV}}$ and δa_{\parallel} . Cases number 5 and 6 have been selected because for both of them δa has a positive impact on the forecast error (i.e. it reduces the forecast error $RE(f^4) < 0$, see Fig. 2) but the impact depends on the evolution of different components. For case 5, the forecast error reduction (see Fig. 3) is due mainly to the evolution of the dropsonde-induced analysis component $\delta a_{\overline{SV}}$ while for case 6 the opposite is true (see discussion in Section 3.2). Furthermore, case 5 can be seen as a typical case with a small and negative projection along the pseudo-inverse ($\rho = -0.05$, see Section 3.3) and case 6 as a typical case with a large positive projection ($\rho = 0.6$). All maps and their corresponding discussion refer to the 500 hPa geopotential height field.

4.1 Case number 5 (18th of February)

On the 18th of February, 17 dropsondes were released in a flight mission from Honolulu. Figure 6a shows the dropsonde-induced analysis difference δa , and Fig. 6c shows the component δa_{SV} in the sub-space spanned by the leading SVs. This comparison shows that δa and δa_{SV} are different ($\psi = 6.4\%$, see Table 1) and δa is characterized by a 5-times deeper structure. It is interesting to compare the error of the 48-hour forecast started from the perturbed initial conditions and valid on the 20th of February at 00GMT. Figure 6 shows the forecast absolute-error difference between $|e^4|$ and $|e^c|$ (Fig 6b), $|e^3|$ and $|e^c|$ (Fig 6d), and $|e^3|$ and $|e^c|$ (Fig 6e). The first thing to note is that the pattern and the intensity of the absolute-error differences shown in Figs. 6b and 6e are very similar and both rather different from Fig. 6d. This confirms the results shown in Fig. 3 that for this case, the evolution of δa and $\delta a_{\overline{SV}}$ has a similar positive impact (i.e. it decreases it) on the forecast error, while δa_{SV} acts to increase it. The impact on the forecast error inside the FVA has been quantified by computing the normalized difference between the root-mean-square error (rmse) [i.e. $(\text{rmse}^j - \text{rmse}^c) / \text{rmse}^c$, for $j = 3, \overline{3}$ and 4], whose value is -9% for $j = 4$ (Fig. 6b), 3% for $j = 3$ (Fig. 6d) and -12% for $j = \overline{3}$ (Fig. 6e).




Figure 7: Case 5, initial state 00GMT of the 18th of February 1998, 500 geopotential height fields. a) δp perturbation, contours every 1.2 m. b) Difference between the 2-day forecast absolute-error of f^4 and f^c started from a_0^4 and C analyses and valid on the 20 February 1998 at 00 UTC. Contour every 10 m, c) δa_{\parallel} perturbation, contours every 0.08 m. d) Difference between the 2-day forecast absolute-error of f^2 and f^c , contour every 1 m. Shaded contours are negative.

Figure 7a shows δp and Fig. 7c shows $\delta a_{||}$. These two initial perturbations are (by construction) identical in shape but have opposite sign and very different magnitude, $\delta a_{||}$ being about 15-times smaller ($\rho=-0.05$, see Table 2). Note that the pseudo-inverse (Fig. 7a) is very different from the dropsonde-induced analysis difference δa (Fig. 6a) and from δa_{SV} (Fig. 6c). Figure 7 shows the absolute-error difference between $|e^1|$ and $|e^c|$, and $|e^2|$ and $|e^c|$. Figure 7b shows that the pseudo-inverse reduces the forecast error over the whole FVA (grey shaded contours) whilst $\delta a_{||}$ slightly increases the forecast error (Fig 7d), in agreement with the fact that ρ is negative and with the RE() results shown in Fig. 4. The normalized difference between the rmse is -35% for f^1 (Fig. 7b) and 2% for f^2 (Fig. 7d).

Figure 8a shows the area of maximum SV concentration, defined as the average of the SV total energy weighted by the amplification factor, and the dropsondes' locations. It can be seen that the dropsondes sample only a small region of the downstream part of the area of maximum SV concentration. For this case, $DLE=2.3\%$ (Fig 5).




Figure 8: Singular vector location (defined as the average total energy weighted by the amplification factor) and dropsondes' locations for (a) case 5 (18 February) and (b) case 6 (20 February).

4.2 Case number 6 (20th of February)

On the 20th of February, 40 dropsondes were released from Hawaii and west of Cape Mendocino. The western flight track was selected by NRL and the eastern track by NCEP. Sondes were deployed on the anticyclonic shear side of the upper level jet, with a good definition of gradients across the lower tropospheric baroclinic zone (*R. H. Langland 2001*, personal communication). Figures 9 and 10 are the equivalent of Figs. 6 and 7 but for this case. Figure 9a shows that δa is characterized by an elongated pattern in the subtropical steering flow, with a first positive maximum centred on the dateline, a dipole structure around 150°W and a final maximum close to the eastern border of the target area T. Figure 9c shows that δa_{SV} is smaller in amplitude than δa (contours are 10 times smaller than in Fig. 9a) with one maximum east of the dateline in correspondence with the first δa maximum and an elongated dipole structure close to the east border of the target area T. Consider now the 48-hour forecast valid on of the 22nd of February at 00GMT. The differences between the absolute-errors $|e^4|-|e^c|$ (Fig.9b) and $|e^3|-|e^c|$ are very similar (Fig. 9d), and both dissimilar to $|e^3|-|e^c|$ (Fig. 9e). The




Figure 9: Case 6, initial state 00GMT of the 20th of February 1998. a) δa perturbation, contours every 5 m. b) Difference between the 2-day forecast absolute-error of f^4 and f^c started from a_0^4 and C analyses and valid on the 22 February 1998 at 00 UTC. Contours every 4 m. c) δa_{sv} perturbation, contours every 0.5 m. d) Difference between the 2-day forecast absolute-error of f^3 and f^c , contours every 4 m. e) Difference between the 2-day forecast absolute-error of f^3 and f^c , contour every 4 m. Shaded contours are negative.

normalized differences of the rmse inside the FVA are -31% for f^4 (Fig. 9b), -22% for f^3 (Fig. 9d) and -8% for f^3 (Fig. 9e).

Figure 10a shows the pseudo-inverse and Fig. 10c shows the analysis component along the pseudo-inverse. The two patterns are identical in shape and sign but have different amplitude, δa_{sv} being about 2-times smaller ($\rho=0.60$, see Table 2). Note that for this case the pseudo-inverse δp (Fig. 10a) and the dropsonde-induced analysis difference δa (Fig. 9a) have both a maximum, east of the dateline, and that δp (Fig. 10a) and δa_{sv} (Fig. 9c) are very similar in shape. Figure 10b shows the difference between $|e^1|$ and $|e^c|$, and Fig. 10d shows the difference between $|e^2|$ and $|e^c|$. These forecast error differences are very similar in shape, with normalized rmse differences inside the FVA of -27% for f^1 (Fig. 10b) and -17% for f^2 (Fig. 10d).

Figure 8b shows the area of maximum SV concentration and the dropsondes' locations. The dropsondes sample one of the two maxima of the SV location. Compared to case 5 (Fig. 8a), there is better agreement between the dropsondes' locations and the area of maximum SV location ($DLE=8\%$, Fig 5).




Figure 10: Case 6, initial state 00GMT of the 20th of February 1998. a) δp perturbation, contours every 1 m. b) Difference between the 2-day forecast absolute-error of f^1 and f^c started from a_0^1 and C analyses and valid on the 22 February 1998 at 00 UTC. Contours every 3 m. c) $\delta a_{||}$ perturbation, contours every 0.5 m. d) Difference between the 2-day forecast absolute-error of f^2 and f^c , contours every 3 m. Shaded contours are negative.

5. Conclusions

Targeted observations are designed to reduce initial uncertainties in the target region T and to reduce the forecast error inside the forecast verification area (FVA). However, mixed forecast results have been obtained from the assimilation of targeted observations during 10 cases of the NORPEX field experiment. Results have in fact indicated that on average the assimilation of targeted data lead to $\sim 2\%$ reduction of the forecast error measured in terms of integrated total-energy, with a peak reduction of 9% (for 2 of the 10 cases). These results cannot be directly compared to the 10% average value obtained by *Szunyogh et al* (2000) and to the 15% obtained by *Montani et al* (1999) because they were based on a single-level fields and not on vertically integrated measures as here. Moreover, *Szunyogh et al* (2000) and *Montani et al* (1999) results refer to 1999 while this study refers to 1998, and the two years are known to be characterized by very different circulation regimes. 1998 was associated with El Niño and characterized by a predominantly zonal flow with a very strong upper level jet; 1999 was ours El Niña year, characterized by a blocked circulation over the West Pacific, a deep trough over Japan and a more pronounced ridge centred on the Pacific.

This paper has investigated possible reasons for the small or negative impact of the targeted observations using a SV-based diagnostic technique. Singular vectors (SVs) identify the phase-space directions along which perturbation growth is maximized during a finite-time interval, and can be used to define a set of diagnostic tools and concepts. For each case, the leading 10 “analysis” SVs, that is, SVs evolving from the analysis time and growing during a 48-h time interval to maximize the total energy norm inside the forecast verification area (FVA), have been computed with a T63L31 resolution model (spectral triangular truncation T63 and 31 vertical levels). The choice of a T63L31 resolution is a compromise between the need of resolving small scales and the limitation of computer usage. The FVA has been set to be 30-60N and 100-130W.

In the first part of this work, the percentage of forecast error explained by a variable number of leading singular vectors has been computed. Results have shown that 44% of the forecast error inside the FVA can be explained by using the 10 leading SVs, and that the use of further 10 SVs only adds a further 3% to this percentage. Following this result, only the leading 10 SVs have been used to define the pseudo-inverse initial perturbation which can correct the most of the forecast error inside the FVA. The fact that the leading 10 SVs define dynamically important directions has been confirmed as the pseudo-inverse initial perturbation when added to the control analysis has always reduced the forecast error (on average

10% when forecast error is measured in terms of vertically integrated total-energy, see also *Buizza et al 1997* and *Gelaro et al 1998*). The pseudo-inverse initial perturbation has been used as a reference in this study.

To investigate the relationship between the dropsonde-induced analysis difference δa , the leading 10 SVs and the pseudo-inverse δp , three initial perturbations have been defined: the dropsonde-induced analysis difference component that belongs to the sub-space defined by the first 10 leading SVs (δa_{SV}), its complement ($\delta a_{\overline{SV}}$) and the δa component along the pseudo-inverse ($\delta a_{||}$). Three indices have been defined to measure the similarity between the five initial perturbations δa , δa_{SV} , $\delta a_{\overline{SV}}$, $\delta a_{||}$ and δp . All these initial perturbations have been defined in terms of the model's vorticity, divergence and temperature and surface pressure fields, thus excluding the humidity field. Changes in the humidity fields due to the assimilation of wind and temperature from dropsondes have been shown not to affect the forecast error in all but case number 3, for which humidity increments were shown to have increased the forecast error by 15%. Once the initial perturbations had been defined, 48-hour forecasts were run from the perturbed initial conditions and the forecast were compared.

Results have shown that on average only 6% of the dropsonde-induced perturbation δa projects onto the sub-space spanned by the leading 10 SVs ($\overline{\psi} \sim 6\%$, see Table 1), with two cases characterized by a 9% maximum projection. In other words, on average 94% of the dropsonde-induced perturbation δa lies in the sub-space orthogonal to the 10 leading SVs. Considering the impact on the forecast error it has been shown that δa_{SV} is dominant in 3 and $\delta a_{\overline{SV}}$ is dominant in 5 out of 10 cases. Moreover, no strong relation has been found between the amplitude of the dropsonde-induced component along the leading singular vectors and the percentage of forecast error variation inside the FVA.

Results have also indicated that in 6 out of 10 cases less than 8% of the dropsonde-induced perturbation projects onto the pseudo-inverse ($\rho < 0.08$, see Table 2), in two cases the projection was $\sim 25\%$ and in two other cases it was $\sim 60\%$ ($0.14 < \rho < 0.60$, see Table 2). Consistently, the two vectors have been almost orthogonal ($\alpha \sim 90^\circ$) in 6 out of 10 cases. In the two cases with the largest projections ($\rho = 0.58$ and $\rho = 0.60$) the forecast error reduction induced by the pseudo-inverse and the dropsonde-induced perturbation have been very similar.

One of the reasons of the small projection of the dropsonde-induced analysis perturbation onto the leading 10 SVs is the limited degree of overlap between the region spanned by the dropsondes and the region of maximum SV concentration. Only case number 6 which is characterized by the largest agreement between the SV and the dropsonde location (Fig 5 and 8) show the closest agreement between the forecast error reduction obtained by correcting the initial condition by the pseudo-inverse and by the dropsonde-induced analysis perturbation (Fig. 4).

In four cases, the pseudo-inverse and the analysis component along it had different sign. In particular, case number 3 with a quite large *DLE* (4%) and ρ (-28%) had opposite forecasts impact because of the opposite sign of the two perturbations.

This can be due to the fact that the effect of the observations on the analysis depends on properties of the assimilation system that are not considered when computing the leading total-energy SVs (e.g. the analysis error covariance matrix which defines the weight the background and the observation have in the analysis). A way to include properties of the data-assimilation system into a SV computation was suggested by *Barkmeijer et al (1999)* who proposed to use an analysis error matrix in the generalized SV computation. *Gelaro et al (2001)* indeed showed that using this norm leads to an increased similarity between the phase-space of the system spanned by data-assimilation and the by the leading SVs during targeted cases, but no conclusions were drawn on the impact on the forecast error.

A promising new way to use analysis error information to define target areas has been proposed by *Baker and Daley (2000)* and *Doerenbecher and Bergot (2001)* and is based on the forecast sensitivity to the observations. Such a technique determines when the forecast is sensitive to the background field or to the observations or to both, avoiding mis-sampling and inefficient use of extra observations. Work along this line should be encouraged.

Acknowledgements

We are very grateful to two anonymous referees for their very helpful comments. We thank Erik Andersson for improving the manuscript. The experimentation was made possible thanks to technical support of Jan Haseler. The figures were skillfully improved by Rob Hine.

Appendix A

A.1 Scalar product and energy norm

Consider the linear space N of vectors \mathbf{x} whose elements x_j are the upper-level vorticity, divergence, temperature and logarithm of surface pressure at different latitude, longitude and vertical coordinates. The total energy norm is defined as

$$\|\mathbf{x}\| = \langle \mathbf{x}, E\mathbf{x} \rangle = \sum_j E_j x_j^2$$

where $E = \text{diag}(E_j)$ is a total energy weight matrix (*Buizza and Palmer 1995*) and x_j is the j -th component of the state vector \mathbf{x} .

A.2 Local projection operator

The local projection operator W is defined as

$$W(\lambda, \varphi)\mathbf{x} = w(\lambda)w(\varphi)\mathbf{x}$$

where λ and φ are the latitude and longitude coordinates, \mathbf{x} is a state vector and $w(\tau)$ is the following weight function

$$\begin{aligned} w(\tau) &= 1 & \tau &= [\tau_1, \tau_2] \\ w(\tau) &= e^{(\tau - \tau_1)/10} & \tau &< \tau_1 \\ w(\tau) &= e^{(\tau - \tau_2)/10} & \tau &> \tau_2 \end{aligned}$$

The local projection operator acts as a smoothed mask. In this work the mask frame is defined by ($\lambda_1=20^\circ\text{N}$, $\lambda_2=60^\circ\text{N}$) and ($\varphi_1=140^\circ\text{E}$, $\varphi_2=240^\circ\text{E}$), and the two couple of coordinate ($\lambda_1=30^\circ\text{N}$, $\lambda_2=60^\circ\text{N}$) and ($\varphi_1=230^\circ\text{E}$, $\varphi_2=260^\circ\text{E}$) define the geographical domain that coincide with the Forecast Verification Area (FVA, see text).

A.3 Singular vectors definition

Let \mathbf{x}_0 be a vector representing a model initial state and \mathbf{x} its 48 hour linear evolution

$$\mathbf{x} = L\mathbf{x}_0$$

L being the tangent model forward propagator. Using the local projection operator W , the total energy norm can be computed inside a specific area (local energy):

$$\langle W\mathbf{x}, EW\mathbf{x} \rangle = \langle WL\mathbf{x}_0, EWL\mathbf{x}_0 \rangle$$

The singular vectors are an orthogonal set of m vectors $\mathbf{v}_i = L\mathbf{v}_i^0$ (orthonormal at the initial time) that maximize the ratio between the final-time local energy norm inside the FVA and the initial total energy norm. The singular vectors \mathbf{v}_i are ordered with decreasing singular value σ_i

$$\sigma_i^2 = \frac{\langle W\mathbf{v}_i, EW\mathbf{v}_i \rangle}{\langle \mathbf{v}_i^0, E\mathbf{v}_i^0 \rangle}$$

In this study, $m=10$ singular vectors are computed using a simplified linear scheme simulating surface drag and vertical diffusion at T63 resolution and 31 model levels.

A.4 Pseudo-initial perturbation

Denote by δe^c the projection of the forecast error e^c onto the first 10 singular vectors,

$$\delta e^c = \sum_{j=1}^{10} \langle W e^c, E W v_j \rangle \frac{v_j}{\sigma_j^2}$$

the pseudo-inverse initial perturbation is defined as the initial perturbation that evolves linearly into δe^c . This perturbation can be written in terms of the initial time singular vectors as follows,

$$\delta p = \sum_{j=1}^{10} \langle W e^c, E W v_j \rangle \frac{v_j^0}{\sigma_j^2}$$

References

- Baker, N. L. and R. Daley, 2000: The observation-targeting problem. *Q. J. R. Meteorol. Soc.*, **126**, 1431-1454.
- Barkmeijer, J., R. Buizza and T. N. Palmer, 1999: 3D-Var Hessian singular vectors and their potential use in the ECMWF Ensemble Prediction System. *Q. J. R. Meteorol. Soc.*, **125**, 2333-2351.
- Bishop, C., and Z. Toth, 1999: Ensemble transformation and adaptive observations. *J. Atmos. Sci.*, **56**, 1748-1765
- Buizza, R., 1994: Sensitivity of Optimal Unstable Structures. *Q. J. R. Meteorol. Soc.*, **120**, 429-451.
- Buizza, R., J. Tribbia, F. Molteni and T. N. Palmer, 1993: Computation of optimal unstable structures for a numerical weather prediction model. *Tellus*, **45A**, 388-407.
- Buizza, R. and T.N. Palmer, 1995: The singular vector structure of the atmospheric global circulation. *J. Atmos. Sci.*, **52**, 1434-1456.
- Buizza, R., R. Gelaro, F. Molteni and T.N. Palmer, 1997: The impact of increased resolution on predictability studies with singular vectors. *Q. J. R. Meteorol. Soc.*, **123**, 1007-1033.
- Buizza, R. and A. Montani, 1999: Targeting observations using singular vectors. *J. Atmos. Sci.*, **56**, 2965-2985.
- Doerenbecher, A., and T. Bergot, 2001: Sensitivity to observations applied to FASTEX cases. *Non-linear Processes in Geophysics*, **8**, 467-481.
- Ehrendorfer, M. and J.J. Tribbia, 1997: Optimal prediction of forecast error covariances using singular vectors. *J. Atmos. Sci.*, **54**, 286-313.
- Emanuel, K., D. Raymond, A. Betts, L. Bosart, C. Bretherton, K. Droegemeier, B. Farrell, M.J. Fritsch, R. Houze, M. Le Mone, D. Lilly, R. Rotunno, M. Shapiro, R. Smith and A. Thorpe, 1995: Report of the first prospectus development team of the U.S. Weather Research Program to NOAA and the NSF. *Bull. Amer. Meteor. Soc.*, **76**, 1194-1208.
- Gelaro, R., R. Buizza, T.N. Palmer and E. Klinker, 1998: Sensitivity analysis of forecast error and the construction of optimal perturbations using singular vectors. *J. Atmos. Sci.*, **55**, 1012-1037.
- Gelaro, R., R.H. Langland, G.D. Rohaly and T.E. Rosmond, 1999: An assessment of the singular-vectors approach to targeted observing using the FASTEX data set. *Q. J. R. Meteorol. Soc.*, **125**, 3299-3327.
- Gelaro, R., T.E. Rosmond and R. Daley, 2001: Singular vector calculations with an analysis error variance metric. To appear in *Mon. Wea. Rev.*
- Gilmour, I, L. A. Smith and R. Buizza, 2001: On the duration of the linear regime: is 24 hour a long time in weather forecasting? *J. Atmos. Sci.*, **58**, 3525-3539.
- Jaubert, G., C. Piriou, S.M. Loehrer, A. Petitpa and J.M. Moore, 1999: Development and quality control of the FASTEX data archive. *Q. J. R. Meteor. Soc.*, **125**, 3165-3188.
- Klinker, E., F. Rabier, G. Kelly and J.-F. Mahfouf, 2000: The ECMWF operational implementation of four-dimensional variational assimilation. III: experimental results and diagnostics with operational configuration. *Q. J. R. Meteorol. Soc.*, **126**, 1191-1218.
- Langland, R.H. and G.D. Rohaly, 1996: Adjoint-based targeting of observations for FASTEX cyclones. Preprints, *7th Mesoscale Processes Conf.*, 9-13 Sept. 1996, Reading, UK., Amer. and Roy. Meteor. Soc., 369-371.
- Langland, R.H., R. Gelaro, G.D. Rohaly and M.A. Shapiro, 1999: Targeted observations in FASTEX: adjoint based targeting procedures and data impact experiments in IOP17 and IOP18. *Q. J. R. Meteorol. Soc.*, **125**, 3241-3270.
- Leutbecker M., J. Barkmeijer, T.N. Palmer and A.J. Thorpe, 2002: Potential improvement to forecasts of two severe storms using targeted observations. *Q. J. R. Meteorol. Soc.*, **128**, 1641-1670
- Mahfouf, J.F., 1999: Influence of physical processes on the tangent linear approximation. *Tellus*, **51A**, 147-166.
- Mahfouf, J.F. and F. Rabier, 2000: The ECMWF operational implementation of four dimensional variational assimilation. Part II: experimental results with improved physics. *Q. J. R. Meteorol. Soc.*, **126**, 564A, 1171-1190.

- Majumdar, S.J., C. Bishop, R. Buizza, and R. Gelaro, 2002: A comparison of ETKF targeting guidance with ECMWF and NRL TE-SVs targeting guidance. *Q. J. R. Meteorol. Soc.*, in print.
- Montani, A., A. J. Thorpe, R. Buizza, and P.Uden, 1999: Forecast skill of the ECMWF model using targeted observations during FASTEX. *Q. J. R. Meteorol. Soc.*, **125**, 3219-3240.
- Palmer, T.N., R. Gelaro, J. Barkmeijer and R. Buizza, 1998: Singular vectors, metrics and adaptive observations. *J. Atmos. Sci.*, **55**, 633-653.
- Pu, Z-X, E. Kalnay, J. Sela and I. Szunyogh, 1997: Sensitivity of forecast errors to initial conditions with a quasi-inverse linear method. *Mon. Wea. Rev.*, **125**, 2479-2503
- Pu, Z-X and E. Kalnay, 1999: Targeting observations with the quasi-inverse linear and adjoint NCEP global models: performance during FASTEX. *Q. J. R. Meteorol. Soc.*, **125**, 3329-3337.
- Rabier, F., E.Klinker, P. Courtier and A.Hollingsworth, 1996: Sensitivity of forecast errors to initial conditions. *Q. J. R. Meteorol. Soc.*, **122**, 121-150.
- Rabier, F., H. Järvinen, E. Klinker, J.-F. Mahfouf and A. Simmons, 2000: The ECMWF operational implementation of four dimensional variational assimilation. Part I: experimental results with simplified physics. *Q. J. R. Meteorol. Soc.*, **126**, 1143-1170.
- Ralph, F. M., Persson, O., Reynolds, D., Neiman, P., Nuss, W., Schmidt, J., Jorgensen, D., King, C., White, A., Bao, J., Neff, W., Kinsmill, D., Miller, D., Toth, Z., & Weilczak, J., 1998: The use of tropospheric profiling in CALJET. *4th Symposium on Tropospheric Profiling: Needs and Technologies*. 20-25 September 1998, Snowmass, CO, 258-260.
- Snyder, C., 1996: Summary of an informal workshop on adaptive observations and FASTEX. *Bull. Amer. Meteorol. Soc.*, **77**, 953-961.
- Thorpe, A. J., and M.A. Shapiro, 1995. *FASTEX: Fronts and Atlantic Storm Track Experiment. The Science Plan*. Available from FASTEX Project Office, July 1995.
- Toth, Z., I. Szunyogh, S. Majumdar, R. Morss, B. Etherton, C. Bishop, S. Lord, M. Ralph, O. Persson and Pu, Z.-X., 2000: Targeted observations at NCEP: toward an operational implementation. *4th Symp. Integrated observing system*, 10-14 Jan. 2000, Long Beach, CA.
- Szunyogh, I., Z.Toth, S. J. Majumdar, R. Morss, C. Bishop and S.Lord 1999: Ensemble-based targeted observations during NORPEX. *3rd Symp. Integrated observing system*, 10-15 Jan. 1999, Dallas, Texas.
- Szunyogh, I., Z.Toth, R.E.Morss, S.J.,Majumdar, B.J. Etherton and C.H. Bishop 2000: The effect of targeted dropsonde observations during the 1999 Winter Storm Reconnaissance Program. *Mon. Wea. Rev.*, **128**, 3520-3537.
- Szunyogh, I., Z.Toth, A.V. Zimin, S.J.,Majumdar and A. Persson, 2002: On the propagation of the effect of targeted observations: The 2000 Winter Storm Reconnaissance Program. *Mon. Wea. Rev.*, **130**, in print.
- Toth, Z., Szunyogh, I., Bishop, C., Majumdar, S., Morss, R., Moskatis, J., Reynolds, D., Weinbrenner, D., Michaud, D., Surgi, N., Ralph, M., Parrish, J., Talbot, J., Pavone, J., and Lord, S., 2002: Adaptive observations at NCEP: past, present and future. Contribution to the *Symposium on observations, data assimilation and probabilistic prediction*, 13-17 January 2002, Orlando, FL.